

Children who continue to have seizures despite medical therapy may be candidates for curative epilepsy surgery



About 1 percent of children ages 17 and under are diagnosed with epilepsy. Medication can control seizures in the majority of patients. However, about 30 percent of children continue to have seizures while on medication. Epilepsy surgery, which was first proposed as a treatment more than a century ago, is highly successful in carefully selected children with medical-refractory seizures, but remains an underutilized option.

UCLA's comprehensive pediatric epilepsy program pioneered the use of state-of-the-art evaluation methods to improve the selection of children, including very young infants, dating back to the mid-1980s and offers multiple surgical and nonsurgical treatment options.

Importance of seizure control

Uncontrolled seizures in pediatric patients can lead to development delay, cognitive or memory deficits, poor quality of life or life-threatening accidents. Children with medically refractory epilepsy should be referred for a diagnostic workup to a high-volume, comprehensive pediatric epilepsy center if they have failed a second medication. Additional attempts at medical treatment after two failed trials of antiepileptics are rarely successful. Surgery is no longer considered an extreme or "last-resort" option for seizure control.

Filling a need to treat uncontrolled seizures

"Pediatric epilepsy patients who fail two or three drugs need alternatives, and we offer many alternatives in our program, both surgical and nonsurgical," says Raman Sankar, MD, PhD, director of the Pediatric Epilepsy Program and chief of pediatric neurology.

"Ongoing seizures are dangerous," says Aria Fallah, MD, assistant professor-in-residence of neurosurgery. "Any child with seizures despite medication therapy needs to be worked up for epilepsy surgery. Epilepsy surgery for a young child may sound radical or scary, but it may be in the best interests of the child. The vast majority of families who go through this say they are extremely grateful and feel like they got their child back.

"Pediatric epilepsy surgery is one of the most remarkable medical advances. We take a chronic, debilitating condition and in many cases are actually able to cure it — and very safely."

Evaluation for surgery

Candidates for epilepsy surgery undergo a full workup to assess the suitability of surgery. At UCLA's dedicated pediatric epilepsy monitoring unit, children are admitted, typically for a few days, for a multi-step evaluation under the direction of a pediatric epileptologist. The workup includes a complete history and physical, video-electroencephalographic (EEG) monitoring and state-of-the-art imaging using 3D MRI and FDG-PET to identify the brain lesions that may be responsible for seizures.

The test results are analyzed during a team conference to determine whether the area of the brain producing seizures can be located and, if so, can be safely excised. While many children can be recommended for surgery on the basis of these results, others may require an additional workup featuring invasive EEG. This is a diagnostic operation in which electrodes are implanted in the brain to more accurately identify the epileptogenic zone and to assess the safety of surgery in that region of the brain.

Excellent surgical outcomes

The development of sophisticated neuroimaging and long-term video-electroencephalographic monitoring has reduced the morbidity and mortality of pediatric epilepsy surgery. Surgical options include hemispherotomy — removal of the entire side of the brain responsible for seizure activity. Preferred for children 2 years old or under, hemispherotomy has been shown to resolve seizures in 80 percent of patients and allows for vastly improved neurological development as the remaining brain takes over the functions of the missing hemisphere. UCLA pioneered the use of hemispherotomy and functional hemispherectomy for select cases of infantile spasms.

More limited surgery can be performed on smaller lesions in the brain responsible for seizures. While this type of operation may involve craniotomy, some patients have the option to undergo an image-guided laser ablation procedure. Most children are discharged from the hospital after one day.

The majority of surgical patients become seizure-free with eventual reduction or sometimes even elimination of drug therapy. Pediatric epilepsy surgery mortality rates are less than 0.5 percent. Children who are not candidates for surgery may benefit from enrollment in a medication clinical trial, vagal-nerve stimulation therapy or a ketogenic (low-carbohydrate and high-fat) diet. UCLA has extensive experience and success in instituting and managing the ketogenic diet as well as the modified Atkins and the low-glycemic-index dietary therapies.

Epilepsy surgery outcomes are highly dependent on physician experience and this surgery is best performed in high-volume centers. Over the past four decades, UCLA physicians have pioneered novel treatments, protocols, diagnostic tools and evidence-based guidelines adopted by many children's hospitals around the world. With more than 800 surgeries performed since the program's inception, UCLA has the experience to offer personalized therapy for individual patients.

A team approach is applied to all cases and includes pediatric neurology, pediatric neurosurgery, pediatric neuroradiology, child psychiatry, pediatric neuropsychology, developmental linguistics and neuropathology.

Mattel Children's Hospital 

Participating Team Members

Gary W. Mathern, MD

Director, Pediatric Epilepsy Surgery Program
Professor, Departments of Neurosurgery and Psychiatry & Biobehavioral Sciences

Raman Sankar, MD, PhD

Director, Pediatric Epilepsy Program
Professor of Neurology and Pediatrics
Rubin Brown Distinguished Chair and Chief of Pediatric Neurology
UCLA Children's Discovery and Innovation Institute

Aria Fallah, MD, MSc

Assistant Professor of Pediatric Neurosurgery

Shaun Hussain, MD, MS

Director, Infantile Spasms Program
Assistant Clinical Professor of Pediatric Neurology

Jason Lerner, MD

Director, Pediatric Neurophysiology Laboratory
Associate Clinical Professor of Pediatric Neurology

Joyce Matsumoto, MD

Director, Dietary Therapies Program
Director, Pediatric Epilepsy Fellowship Training Program
Assistant Clinical Professor of Pediatric Neurology

Lekha Rao, MD

Assistant Clinical Professor of Pediatric Neurology

Joyce Wu, MD

Director, Tuberous Sclerosis Program
Clinical Professor of Pediatric Neurology

Sue Yudovin, RN, MN, CPNP

Patient Care Coordinator, Epilepsy Surgery Program

Kristina Murata, NP

Patient Care Coordinator, Ketogenic Diet Clinic

Contact Information

Department of Neurosurgery
Edie & Lew Wasserman Building
300 Stein Plaza, Suite 420
Los Angeles, CA 90095

(310) 825-5111

Division of Pediatric Neurology
UCLA Children's Health Center
200 UCLA Medical Plaza, Suite 265
Los Angeles, CA 90095

(310) 825-0867 Appointments and referrals

(310) 825-6196 Administrative office and information

neurosurgery.ucla.edu/pediatric-epilepsy