

SOLID TUMOR ANCILLARY TESTING

BREAST SERVICE:

NOTE: For all metastatic breast cases, please give the case to breast service (WW breast service or SM breast attendings) to sign out the biomarkers. The case can be signed out by the original service (according to organ), and then give to breast service to put an addendum for biomarkers.

Primary Breast	Immunohistochemistry	FISH
All primary carcinomas (including metaplastic)	ER, PR, HER2, Ki-67	HER2
Ductal carcinoma in situ (DCIS)*	ER, PR*	
Metastatic Breast	Immunohistochemistry	FISH
All breast carcinomas (including metaplastic)	ER, PR, HER2, Ki-67	HER2

*Performed only on resection/excision specimen

- Smart text: .breastbiomarkers

GASTROINTESTINAL SERVICE

Primary Colorectal	Immunohistochemistry	FISH	Molecular
Adenocarcinoma	MMR (does not need to be repeated if performed on initial biopsy specimen)		MSI ¹ , BRAF ²
Adenocarcinoma (Biopsy)	MMR (Perform on all cases if adequate tissue present)		
Metastatic Colorectal	Immunohistochemistry	FISH	Molecular
Adenocarcinoma	MMR ¹		KRAS*

Primary Pancreas	Immunohistochemistry	FISH	Molecular
Ductal adenocarcinoma	Smad4 (Performed on resection specimen)		
Metastatic Pancreaticobiliary	Immunohistochemistry	FISH	Molecular
Metastatic pancreatic ductal adenocarcinoma	MMR ¹		
Metastatic cholangiocarcinoma	MMR ¹		

GI Stromal Tumor & Neuroendocrine	Immunohistochemistry	FISH	Molecular
GI neuroendocrine tumor (all sites, including metastasis)	Ki-67		
Primary GIST (all sites)			PDGFRA/c-kit

Primary Gastric and GEJ	Immunohistochemistry	FISH	Molecular
Adenocarcinoma	HER2, PD-L1 [^]	HER2 (only if IHC is 2+)	
Metastatic Gastric and GEJ	Immunohistochemistry	FISH	Molecular
Adenocarcinoma	HER2, MMR ¹	HER2 (only if IHC is 2+)	

Primary Esophageal	Immunohistochemistry	FISH	Molecular
Adenocarcinoma	HER2 (clinician request)	HER2 (only if IHC is 2+)	

<u>Primary Ampulla</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
Adenocarcinoma (resection)	CK7, CK20, CDX2, MUC1 and MUC2		

*If KRAS result is wild type, order reflex CRC Panel sequencing analysis (BRAF, KRAS, NRAS, PIK3CA, and AKT1). In addition, order reflex HER2 IHC. Order HER2 FISH if IHC score is 2+.

¹ All metastatic GI malignancies (including pancreaticobiliary malignancies) are being tested for MMR deficiency by IHC given the approval of PD1 inhibitors in all GI cancers that are MSI-H

¹ MSI by PCR is indicated if MMR IHC results are equivocal or questionable. If IHC has already been done on a biopsy and a normal expression pattern is observed, MSI PCR will not be performed on the resection specimen from the same patient **UNLESS:**

- 1) Patient age under 50
- 2) Personal hx of Lynch-related tumor(s) – may need to be informed by clinicians
- 3) Family hx of CRC or Lynch syndrome – may also need to be informed by clinicians
- 4) Histologic features suggestive of MSI on resection specimens (mucinous, poorly differentiated, medullary, tumor infiltrating lymphocytes, Crohn-like peritumoral lymphoid response)

² BRAF mutational analysis is indicated for cases with loss of MLH1 expression.

[^] Use the PD-L1 stain Smart text: .PDLGEJ, can be used for all GI PD-L1 IHC reports (CPS)

c-KIT/PDGFR is a send out test. Please email SurgicalPathologySendouts@mednet.ucla.edu to order the test (see below).

LUNG SERVICE

<u>Primary Lung</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
Adenocarcinoma	PD-L1	Lung panel	Lung panel
Squamous cell CA	PD-L1		
Small cell carcinoma or carcinoid	PD-L1 (Clinician request)		
Mesothelioma	PD-L1, BAP1 if requested		
<u>Metastatic Lung</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
All carcinomas	PD-L1 (Clinician request)	per request	per request

- Smart text: .IMPDL1

GENITOURINARY SERVICE

<u>Metastatic Tumors</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
Urothelial carcinoma	PD-L1		
<u>Locally advanced tumors</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
Urothelial carcinoma	PD-L1 (per clinician request)		

DERMATOLOGY SERVICE

<u>Tumor</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
Melanoma			BRAF (per request)
Metastatic melanoma			BRAF (per request)

GYN SERVICE

<u>Primary Ovarian</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
Endometrioid and clear cell and other uterine carcinomas	MMR*, ER, PR, p53	
Uterine high grade serous carcinoma	p53, WT1, p16, ER, PR	
Most high grade carcinomas	p53, WT1, p16, ER, PR	
<u>Metastatic Ovarian</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
Endometrioid and clear cell type carcinomas	MMR, ER, PR (per clinician request and if not performed on primary)	
All high grade carcinomas (includes MMT)	p53, ER, PR (per clinician request and if not performed on primary)	

<u>Primary Uterine</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
All endometrial carcinomas	MMR, ER, PR	
<u>Metastatic Uterine</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
All endometrial carcinomas	MMR, ER, PR (per clinician request and if not performed on primary)	
All high grade carcinomas (includes MMT)	HER2, p53 (per clinician request and if not performed on primary)	HER2 (per clinician)

<u>Primary Cervical</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
Squamous cell carcinoma	P16, Ki-67, and HPV ISH (if needed)	
Adenocarcinoma	P16, Ki-67, and HPV ISH (if needed)	
<u>Metastatic Cervical</u>	<u>Immunohistochemistry</u>	<u>FISH</u>
Squamous cell carcinoma	P16, Ki-67, and HPV ISH (if needed)	
Adenocarcinoma	P16, Ki-67, and HPV ISH (if needed)	

*Performed on excision/resection only, unless requested by clinician to be performed on initial biopsy specimen.

- MLH1 promoter hypermethylation studies are ordered for cases with loss of MLH1 expression. Please email SurgicalPathologySendouts@mednet.ucla.edu to order the test (see below).

NEUROPATHOLOGY SERVICE

<u>Tumor</u>	<u>Immunohistochemistry</u>	<u>FISH</u>	<u>Molecular</u>
GBM (Grade IV)	GFAP, Ki67, IDH1 R132H (Add Olig2 if concerned for PNET or ependymoma)	PTEN, EGFR	MGMT methylation IDH1 #
Infiltrating glioma (Grade II, III)	GFAP, Ki67, IDH1 R132H, ATRX, p53	1p/19q	MGMT methylation
Midline glioma (Infiltrating glioma in thalamus, brainstem, spinal cord)	H3K27M (in addition to infiltrating glioma workup, see above)		
Pilocytic astrocytoma	BRAF V600E, GFAP, Ki-67, neurofilament, synaptophysin, IDH1 R132H	BRAF	
Ganglioglioma	BRAF V600E, GFAP, Ki-67, neurofilament, synaptophysin		
Pleomorphic xanthoastrocytoma	BRAF V600E, GFAP, Ki-67, neurofilament, synaptophysin		
Pituitary adenoma	IM PIT panel (LH, FSH, TSH, GH, prolactin, ACTH, Ki-67) *		
Chordoma	S100, Ker AE1/3, Brachury		
Meningioma vs Solitary Fibrous Tumor/Hemangiopericytoma	EMA, SSTR2A, STAT6, Ki67 (Classic or usual meningioma does not require these stains)		
Medulloblastoma ^	Beta catenin, synaptophysin, GFAP, Ki67, YAP1, GAB1	N-MYC, C-MYC	
Ependymoma	GFAP, Ki67 if classic (add Synaptophysin, neurofilament, EMA, CD99, D2-40, Olig2 if distinguishing from astrocytic tumors)		

if IDH1 IHC is negative and patient is less than 54 years old, order IDH1/IDH2 PCR unless Foundation Medicine genomic profiling is ordered.

*PIT1, SF1, synaptophysin IHC if hormone stains from the initial hormone panel are negative

^ If nodular/desmoplastic variant suspected add reticulin special stain.

^Add INI1 IHC if AT/RT is suspected or less than 10 years old; Foundation Medicine genomic profiling via Neurooncology recommended.

MGMT methylation studies is a send out test. Please email SurgicalPathologySendouts@mednet.ucla.edu to order the test (see below).

HEAD AND NECK SERVICE: p16/HPV ISH staining flow chart




Figure 1. High-risk human papillomavirus (HR-HPV) testing in head and neck squamous cell carcinomas (SCCs). Abbreviations: IHC, immunohistochemistry; OP, oropharyngeal. ¹Consider HR-HPV-specific testing for equivocal p16 results (50%–70% nuclear and cytoplasmic staining). ²May also be reported as p16 negative with a comment specifying that the tumor is very likely HPV negative. ³May also be reported as p16 positive with a comment specifying that the tumor is very likely HPV positive. ⁴HR-HPV may be indicated in patients where the clinical suspicion for an HPV-positive SCC is high. ⁵Consider Epstein-Barr encoding region (EBER) in situ hybridization for Epstein-Barr virus for the rare metastatic nonkeratinizing squamous cell carcinoma that is HR-HPV negative. ⁶Include comment, "Likely oropharyngeal primary."

ANCILLARY SEND OUT TESTS:

<u>Test Name</u>	<u>Department</u>	<u>Location</u>
MGMT (Molecular)	NP	NeoGenomics
BRAF V600E (Molecular)	NP	NeoGenomics
BRAF Rearrangement FISH for PA	NP	NeoGenomics
MLH1 Promoter Methylation (Molecular)	GYN	NeoGenomics
cKIT w/reflex to PDGFRa	GI/BST	Knight Diagnostic Laboratories/OHSU
cKIT for Melanoma	GI/Derm	Knight Diagnostic Laboratories/OHSU
Gene Trails Panel for Hematologic Malignancies	HemePath	Knight Diagnostic Laboratories/OHSU
Iron Quant	Liver	Mayo
Copper Quant	Liver	Mayo
Amyloid Protein ID	H/L, GI/Liver	Mayo
PLA2R IF	Renal	Mayo
Alport Staining for Collagen IV	Renal	Dr. Laura Flynn, Seattle Children's
AFB, Bacterial, Fungal Broad-range PCR	All	Univ. of Washington
MAML2 FISH	Head/Neck	Mayo
MYD88	HemePath	NeoGenomics

For all send out tests please e-mail SurgicalPathologySendouts@mednet.ucla.edu with the required test, case number and case block. Please check with them if a circled H&E slide is required.