

Orthopaedic Total Joint Replacement Pre-Op Class

Presented by UCLA Nursing, Pharmacy, Physical Therapy, and Occupational Therapy

Welcome!





Class Outline

- I. Nursing
- II. Pharmacy
- III. Physical Therapy
- IV. Rehab Demonstration
- V. Occupational Therapy
- VI. Ortho Nurse Navigator/Case Management



Review these electronic attachments


- Surgeon's Discharge Instructions
- Class Powerpoint
- Knee Rehab Brochure
- Knee Exercise Program
- Physical Therapy Locations
- Pain Scale Chart
- Local Equipment Stores List
- Fall Prevention Tips
- Intimacy after Joint Replacement





NURSING

Your Surgery



Arrival Time to the Hospital

You will receive a phone call from UCLA Santa Monica

1 Business Day

prior to your surgical date, alerting you of your arrival time to the hospital.

- Call can come between (9am to 6pm)
- •If you do not receive a call 1 business day before your surgery, then please call (424) 259-6337 after 3pm.

If still no call after 6pm, please call Ortho Nurse Navigator, Wilson Phoeng at (310) 295-7403



Day Before & Day of Surgery

Patient Checklist - Your Diet

No food or drink after instructed time

Arrival Time	Stop Solid Foods	Drink Clear Liquids Until
6 a.m.	10 p.m.	4 a.m.
8 a.m.	Midnight	6 a.m.
10 a.m.	2 a.m.	8 a.m.
12 p.m.	4 a.m.	10 a.m.
2 p.m.	6 a.m.	12 p.m.

Clear, see-through liquids include:

+Water +Pedialyte +Gatorade +Powerade

Clear Liquids are encouraged until 2 hours before your scheduled arrival time at the hospital



Day Before & Day of Surgery

Patient Checklist – Medication & Hygiene

Medications

Take any Blood Pressure medications with small sips of water

Hygiene

- Shower and wash your hair
- You will be given a special liquid soap prior to your surgery during you surgeon's clinic.
- + Hibiclens Instructions
- Do not shave at or near the site of surgery
- Remove fingernail polish and acrylic nails

Risk for Infection



Report any changes in physical condition (cold, fever, rash, nausea vomiting, diarrhea)



Day Before & Day of Surgery

Patient Checklist – What to Bring & What Stay at Home

What to Bring With You


- List of meds and supplements
- Copy of Medical & Surgical Hx
- Insurance card, Driver's license
- Co-pay (check with insurance)
- Copy of Advanced Directive
- Eyeglasses, Hearing Aid w/ battery
- Important telephone numbers
- Payment for discharge medications credit card
- CPAP, rescue inhalers

What To Leave At Home

- Jewelry (Wedding bands OK)
- Checkbooks
- Common Medications
 (If unsure, please check with Ortho Nurse
 Navigator to ensure that we carry your particular medications here at UCLA)



Day of Surgery





PACU – Same Day Discharge

Unit Overview

Patients

Surgeries: Knee, Hip, Spine, Shoulder

Length of Stay: 0 to 2 days

26 Private Rooms

Staff

- Primary Nursing Team: RN & CCP
- · Support: Charge Nurse, Management
 - Ancillary Staff

Visiting Hours

https://www.uclahealth.org/covid-19-visitor-restrictions



3NW Spine & Ortho

Unit Overview

Patients

Surgeries: Knee, Hip, Spine, Shoulder

Length of Stay: 0 to 2 days

26 Private Rooms

Staff

- Primary Nursing Team: RN & CCP
- · Support: Charge Nurse, Management
 - Ancillary Staff

Visiting Hours

https://www.uclahealth.org/covid-19-visitor-restrictions



Foods, Fluids, and Elimination

Management of Issues

- Nausea
- Constipation

Progression

- IV hydration
- Clears to Regular

Importance

- Have energy, heal better
- Good nutrition, better elimination
- Stay hydrated





Mobility & Safety

 Call for assistance before getting up Everyone is a Fall Risk!!

(Medications, Lines, Pumps, and Orthostatic Hypotension)


- Do not feel like calling = bothering
- You can ambulate with any staff member



Deep Vein Thrombosis (DVT) Prevention

Purpose

- Better circulation prevent blood pooling
 - Prevent Blood Clots/PE
 - Prevent Complications (i.e., Stroke)



Tools/Interventions

- Sequential Compression Devices (SCDs) or Foot Pumps
 - Blood thinners
 - Ankle pumps
 - Move more!





Infection Prevention (Continued)

Purpose

- Prevent complications
- Heal better
- Prevent readmissions

Interventions

- Post-op IV antibiotic
- Hand washing
- Dressing change

What to Watch Out For

Redness, Swelling, Drainage, Fever





Swelling Management

To help reduce swelling on operated leg

- Change positions every hour
- Sit for less than 1 hour at a time
 - · Out of bed to chair for meals

Elevated affected extremity above the heart

 If increased swelling is noted in foot or ankle, it is time to spend more time with the leg elevated

"Toes above the Nose"

- · Use ice, but do not get the surgical dressing wet
- Ankle Pumps while elevating extremity above the heart

Continue for 6-8 hours/day for 6-8 weeks after surgery





TKA Wedge Pillow (aka "LRU Pillow")



Foam wedge pillow placed under surgical leg for Total Knee Arthroplasty (TKA) patients after surgery

TKA Wedge Pillow (Continued)

TKA Wedge Pillow-Instructions:

- Rest your leg on the pillow at all times when not moving x 4 weeks (May remove for walking, toileting, eating, etc...)
- Sleep on TKA pillow as much as possible, <u>unless</u> significantly interfering with sleep

Recommendations/Considerations:

- Place a towel on top of pillow to keep it clean
- Pad your heel with cushioned socks or bandage if irritation occurs
- You can use the TKA pillow on all surfaces
- You can and should use ice on your knee when on the pillow
- After 4 weeks, you may reduce use to "as needed"
- Continue use if significant knee swelling persists
- Continue use after a lot of exercise or activity to prevent swelling from returning







TKA Wedge Pillow - Ordering Instructions

(You are encouraged to purchase this item <u>2 weeks before surgery</u> to ensure immediate use at home after surgery)

1) Go to www.MSMProductsIlc.com

> Products > LRU Pillow



2) Scroll down the page and "click" the red box



3) Pick "<u>High Density Foam</u>" (\$99) for Uncovered Pillow

"Vinyl Covered Pillow" (\$149)

> Add to Cart- then check out!





Discharge Day – Post-Op Day 0-2

MD written order is a must! /ill



- Discharge time goal 11:00am
- Arrange transportation



- Discharge instructions & Follow-Up
 - Activities: Showering, Driving
 - Incision site care/Dressing change
 - Follow-up appointments
 - When to call your doctor
 - No formal therapy, Outpatient or Home Health





⚠(Bring payment for medication co-pay. Have credit card payment ready on day of discharge for medication co-pay)



PHARMACY

Medication Considerations **BEFORE** Surgery

Inform physician of medication allergies of adverse reactions

Review your current medications

- · Prescription drugs
- Over-the-counter (OTC) medications
- Vitamins
- Herbal Supplements

Medications to avoid or stop prior to surgery

- Blood thinners should be stopped
- Most medications can be taken up until the night before surgery
- · Consult your physician about timing

Use of personal medications

- Not allowed
- Alternatives on formulary can be ordered. (Consult the physician or pharmacist)





Medication Considerations <u>AFTER</u> Surgery

Antibiotics

Used to prevent infections at the surgical site

- ✓ Cefazolin (Ancef) and vancomycin are commonly used post-operatively
- ✓ Usually given as a one-time dose hours after the end of surgery

Pain Control

Various drugs can be used to prevent or treat pain

These may include, but are not limited to:

- ü Hydrocodone/acetaminophen (Norco, Vicodin), Oxycodone
- ü Tramadol, Tylenol, Celebrex
- ü Hydromorphone, morphine
- ü On-Q Pump Peripheral nerve block for knee replacement called Ropivcaine



Medication Considerations <u>AFTER</u> Surgery

No invasive procedures including dental work for 3 months after date of surgery

❖ Unless emergency or infection concern – please notify your joint surgeon immediately

After 3 months from surgery you must take preventive antibiotics for all dental work and invasive procedure

Procedure	Antimicrobial Agent	Dose	Timing	Duration
Dental	cephalexin, cephradine, amoxicillin	2 gm PO	1 hr prior to procedure	Discontinued within 24 hrs of procedure
Ophthalmic	gentamicin, tobramycin, ciprofloxacin, gatifloxacin, levofloxacin, moxifloxacin, ofloxacin, or meomycin-gramicdin-polymyxin B cefazolin	Multiple drops topically over 2 to 24 hrs or 100 mg subconjunctivally	Consult opthamologist or pharmacist for dosing regimen	Discontinued within 24 hrs of procedure
Orthopaedic	cefazolin cefuroxime or vancomycin	1-2 g IV 1.5 g IV 1 g IV	Begin dose 60 min prior to procedure	Discontinued within 24 hrs of procedure
Vascular	cefazolin or vancomycin	1-2 g IV 1 g IV	Begin dose 60 min prior to procedure	Discontinued within 24 hrs of procedure
Gastrointestinal Esophageal, Gastroduodenal Biliary Tract Colorectal	cefazolin cefazolin neomycin + erythromycin base (oral) or metronidazole (oral)	1-2 g IV 1-2 g IV 1 g 1 g	Begin dose 60 min prior to procedure Dependent on time of procedure consult with Physican/Pharmacist	Discontinued within 24 hrs of procedure
Head and Neck	clindamycin + gentamicin or cefazolin	600-900mg IV 1.5 mg/kg IV 1-2 g IV	Begin dose 60 min prior to procedure	Discontinued within 24 hrs of procedure
Obstetric and Gynecological	cefoxitin, cefazolin ampicillin/sulbactam	1-2 g IV 3 g IV	Begin dose 60 min prior to procedure	Discontinued within 24 hrs of procedure
Genitourinary	ciprofloxacin	500 mg PO or 400 mg IV	1 hr prior to procedure Begin dose 60 min prior to procedure	Discontinued within 24 hrs of procedure



Medication Considerations <u>AFTER</u> Surgery (Continued)

Various medications are used to prevent and treat:

Nausea and Vomiting

Zofran® (ondansetron), Reglan® (metoclopramide)

Constipation

• Colace® (docusate), Miralax® (polyethylene glycol), Dulcolax® (bisacodyl), Fleet's enemas®

Insomnia

Ambien® (zolpidem), Restoril® (temazepam)

Acid Reflux/Indigestion

• Pepcid® (famotidine), Protonix® (pantoprazole), Maalox®



Medication Considerations AFTER Surgery (Continued)

Formation of blood clots can be a consequence of your surgery

Blood thinners are given to **prevent clots**

Subcutaneous Injection

- Lovenox® (enoxaparin)
- Heparin
- Arixtra® (fondaparinux)

Oral

- Aspirin
- Coumadin® (warfarin)

Inform your physician if you were taking these prior to surgery or if you have a history of bleeding (such as GI bleeding, blood in the urine, or "easy" bruising)



PHYSICAL THEBAPY



Rehabilitation – Physical Therapy

Post-Operative Day #0

- PT Begins <u>day of</u> surgery for Primary Total Knee Replacements
 - Evaluation occurs in the recovery unit, especially if plans for possible same day discharge
 - PT evaluation is around 90 minutes after surgery or after resolution of surgery side effects
 - Focus on bed mobility, transfers, and household level ambulation
 - Stair training to also occur on an individualized basis
 - May be cleared for same day discharge after PT, provided MD and medical clearance
 - · If staying overnight, evaluation may occur on 3NW unit instead
 - Exceptions:
 - Latest scheduled cases (or those with significant unplanned delays) will be seen the following morning POD #1 by PT on the Ortho unit, but can participate in early mobility with RN the night of surgery
 - Total Knee Resections or Revisions will not see PT until POD #1



Rehabilitation – Physical Therapy

Post-Operative Day #1

- Patients staying overnight will have PT Evaluation and/or Treatment the morning of POD #1
- PT frequency-Primary Total Knee Replacement
 - 1st visit am of POD #1
 - Focus on bed mobility, transfers, and household level ambulation. Review of exercises.
 - Stair training to also occur on an individualized basis.
 - DC from PT after am session if goals are met
 - If staying longer for medical reasons and/or goals are not met, 2nd session to occur in the afternoon

**PT frequency-Total Knee Resection or Revisions to be seen 1-2 times per day during length of stay until discharge or until goals met, to be determined based on individualized needs and medical complexity



Daily Physical Therapy Treatments

Primary focus is on safe discharge

- Progressing safe household level mobility and walking
- Increasing Knee Range of Motion
- Stair/Step training (as needed)
- Car transfer training
- Caregiver/"Coach" Training





Daily Physical Therapy Treatments Con't.

- Ongoing pain management is key!
 - Important for optimal progression during PT sessions
 - Ideal pain level is 4 or less (on a 1-10 scale) during activity
 - Be sure to communicate your pain level to your nurse with movement of the surgical leg
 - Alternatives to pain medicine
 - Mobilize-"Movement is medicine!"
 - Cold packs or ice
 - Positioning



Coordinating Your Care

- To get the most out of your session, you can best prepare by:
 - Eating meals early (7-8am)
 - Performing hygiene and toileting needs prior to therapy
 - Defer all phone calls and/or visitors before/after therapy
 - Exception: You are encouraged to have one visitor who will be your "coach" at discharge to participate in therapy sessions
 - Manage your pain with RN
 - Take oral medications on schedule




Movement Precautions – Knees

Knee Precautions

- Avoid twisting on the operative leg when standing or walking
 - Take small steps to turn
- Avoid putting pillows directly behind the knee. Keep it straight in bed!
- Prevent swelling
 - Change positions frequently
 - Elevate your leg in-between activities such as standing, walking, and exercises







Questions?

- Up next
 - Practice sessions with PT/OT
 - Getting out of bed
 - Walking with a Walker
 - Exercises
 - Activities of daily living



OCCUPATIONAL THERAPY



Rehabilitation – Occupational Therapy

Post-Operative Day #0 or 1

- Occupational Therapy
 - POD #0: Cases pre-determined to be discharging same day may be seen by OT along with PT
 - If staying overnight, OT begins POD #1
 - DC from OT after evaluation if goals are met
 - Focus on everyday activities
 - Getting dressed
 - Moving in the bathroom
 - Home Activities



Preparing for Home

Preparation is key! **Prior** to surgery, you should:

- Have an adequate support system in place to help with preparing meals, bathing, and moving around the house
- Plan on having a designated person/s physically able to help you upon discharge. This can be:
 - Family and/or close friends
 - Private caregiver if needed
 - Interview and hire ahead of time.
 - Housekeeper, gardener, etc.
 - Arrange in advance as needed



Preparing for Home

Is your home set up?


- You should have, or be able to obtain, all the *durable medical equipment* necessary, such as:
 - Front wheeled walker
 - Commode or raised toilet seat
 - Not required, but may ease transfers for taller patients or those with co-morbidities struggling to rise from low surfaces
- Prepare your home ahead of time to accommodate a walker and prevent falls (see handouts)
- Set up bedroom accommodations on the **first floor** (if you live in a 2 story home)



Preparing for Home – Common Bathroom Equipment


- Raised toilet seat/Commode
- Grab Bars
- Shower seat
- Tub Transfer Bench
- Non-skid Bath Mat/Strips
- Hand Held Shower Head







Preparing for Home – Common Home Modifications





Discharge Planning to Home

- We hope to prepare you for a safe return home on the same day or 1st day after surgery.
 - With a little assistance or supervision, you should be able to:
 - Get in and out of bed
 - Walk household distances with a walker
 - Perform steps/stairs with the appropriate assistive device
 - Use the toilet or commode
 - Complete self-care activities



CASE MANAGEMENT

3 KEYS to Knee Recovery



- 1. Perform home knee exercises every day 3 times a day.
 - ✓ Have formal physical therapy 2 to 3 times a week for 6 to 8 weeks
 - ✓ Range of Motion Goal: 0 degree extension (knee completely straight) to 120 degree flexion (knee bend) by 6-8 weeks after surgery.



2. Manage knee swelling:

- ✓ Elevate operated knee toes above the nose for 6 to 8 hour per 24 hours
- ✓ Ice 4 to 5 times per day 20-30 mins each time



3. Manage your pain:

- ✓ Take your pain medication as needed to get you through the exercise, walking, and physical therapy.
- ✓ As pain gets better, start weaning off the stronger pain medications first.

Designated Helper – Family/Friends

Prior to surgery, designate one or more helpers to assist you after surgery for 1 to 2 weeks.

(Time frame will <u>vary</u> depending on your recovery progress after surgery)

Designated Helper's possible duties:

- · Cooking, cleaning, laundry.
- Assisting you with navigating your home safely.
- · Driving you to your appointments.
- Encouraging you to perform your exercises safely.





Discharge Planning

- Home → No formal therapy or Outpatient physical therapy (after cleared by MD)
- 2. Home with Home Health services (RN & PT/OT) → Outpatient physical therapy (after cleared by MD)
- 3. Skilled Nursing Facility (SNF) → Home
 - Santa Monica Health Care Center UCLA Health designated Center of Excellence for Joint Replacement
 - 1320 20th St, Santa Monica, CA 90404
 - Phone (310) 829-4301



Discharge Planning Con't.

Very **few** patients need SNF rehab before they are safe to go home

- Need maximal or 2 person assistance
- Cannot walk or take steps
- Frequently losing your balance
- ⚠•Medical issues that require skilled nursing need ⚠



Medicare Part A & B

Removal of Total Knee Replacement from Inpatient Only List

- Change was made on Jan 1, 2018 for total knee replacement
- All procedures will be done as outpatient procedure, unless there is medical justification to change to inpatient.
- Cannot discharge to SNF unless made inpatient with medical justification.
- Possible Co-pay of up to 20% for outpatient procedure
 - Typically covered by secondary insurance.

Durable Medical Equipment

Medicare Covered Items

- One item within the last 5 years
- Walker or ambulation device



Self-Pay Items

(Commode, Raised Toilet Seat, Shower Chair, Adaptive Equipment)

- Local equipment stores (see handout)
- Online/Amazon
- Hospital DME vendor
- *Non-Medicare patients: Check your insurance provider about specific DME coverage



Home Health Services

- Agency will call you to make home visit arrangements
- Nursing
 - · RN will visit your home within 24-48 hours after discharge
- Physical Therapy/Occupational Therapy (1-3 times per week)
 - Evaluation by home therapist
- Home health PT/OT discontinued prior to outpatient therapy
- Check with your insurance provider about your Home Health coverage



Skilled Nursing Facility (SNF)

- Short-term rehabilitation
 - Therapy 1-3 hours per day
- Transportation to SNF may NOT be covered
- Your case manager will arrange admission and transportation
- Check with your insurance provider about your specific coverage
 - May be dependent on the hospital assessment of needs and progress


Private Pay Caregiving Agencies

24 hour Private Caregiving Service at home

- 24 Hr Home Care: 310-258-9525
- Accredited Home Care: 800-974-1234 ext 205
- 123 Home Care: 310-361-8558
 - Average cost \$30-40 dollars an hour please call for rates.
 - You can hire same day from 4 hours to 24 hours.
 - You can interview staff ahead of time.



Class Survey – See instructions below!





Orthopaedic Nurse Navigator

- For any orthopaedic related questions and concerns, please contact Orthopaedic Nurse Navigator
- Contact Information:
 - Wilson Phoeng RN, MSN, ONC, PHN
 - Email: wphoeng@mednet.ucla.edu
 - Office # 424-259-8266
 - Work Cell # 310-295-7403
 - Pager # 800-233-7231
 - Page ID # 96637





Thank You All For Coming





UCLA Health