

# NTOS-Q

## Neurogenic Thoracic Outlet Syndrome Questionnaire

Please answer the following questions based on your symptoms during the past 30 days. For each item, check the box that best represents your experience.

### Section A: Symptom Severity

Check one box from 0 (None) to 10 (Worst imaginable)

1. Pain in the neck or supraclavicular region

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

2. Pain in the shoulder, arm, or hand

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

3. Tingling or numbness in the arm or hand

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

4. Weakness or heaviness in the arm or hand

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

5. Symptoms worsening with overhead activity

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

6. Symptoms worsening with prolonged use (e.g., typing, driving)

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

### Section B: Functional Limitations

0 = No limitation, 10 = Completely unable

7. Personal care (e.g., dressing, washing)

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

8. Household tasks (e.g., cooking, cleaning)

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

9. Work or school activities

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

10. Recreational/sports/hobby activities

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

11. Carrying or lifting objects

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

12. Using hands for fine motor tasks (e.g., typing, writing, buttoning)

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

## Section C: Impact on Quality of Life

0 = No impact, 10 = Severe impact

13. Sleep quality

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

14. Anxiety or depression related to NTOS

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

15. Social impact (e.g., limiting time with friends/family)

0 ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10 ☐

## Section D: Overall Impression

16. How would you rate your overall NTOS symptoms today? (Check one)

- ☐ None
- ☐ Mild
- ☐ Moderate
- ☐ Severe
- ☐ Very Severe

**Name:**

**DOB:**